



# flashBAC™ Compatible Vectors

This table provides a summary of different baculovirus transfer vectors that have been produced by many laboratories over the past 30 years. These include vectors with polyhedrin promoters, p10 promoters, p6.9 promoters and gp64 promoters. Although the complete sequences of some of these vectors are not available we have indicated the restriction enzyme cloning sites. The vectors listed are all compatible with flashBAC™ or BacPAK6.

Type (locus)	Vector (Size)	Promoter(s)	Features within Cloning Sites	Additional Features/Notes
Single Promoter (polyhedrin)	pVL1392 (9.6 kbp)	polyhedrin	Bgl II, Pst I, Not I, Eag I, EcoR I, Xba I, Sma I, BamH I	
	pVL1393 (9.6kbp)	polyhedrin	BamH I, Sma I, Xba I, EcoR I, Not I, Eag I, Pst I, Bgl II	
	pAcG1 (8.5 kbp)	polyhedrin	ATG, GST, BamH I, Sma I, EcoR I	
	pAcG2 (8.5 kbp)	polyhedrin	ATG, GST, Thrombin cleavage-BamH I, Sma I, EcoR I	
	pAcG3X (8.5 kbp)	polyhedrin	ATG, GST, Factor Xa cleavage, BamH I, Sma I, EcoR I	
	pAcGHLT-A (8.7 kbp)	polyhedrin	ATG, GST, BamH I-6x His tag, Protein kinase A site, Thrombin cleavage-Nde I, EcoR I, Stu I, Nco I, Sac I, Not I, Sse8387 I, Pst I, Kpn I, Sma I, Bgl II	

	pAcGHLT-B (8.7 kbp)	<i>polyhedrin</i>	ATG, GST, <i>BamH I</i> , 6x His tag-Protein kinase A site, Thrombin cleavage- <i>Xho I</i> , <i>EcoR I</i> , <i>Stu I</i> , <i>Nco I</i> , <i>Sac I</i> , <i>Not I</i> , <i>Sse8387 I</i> , <i>Pst I</i> , <i>Kpn I</i> , <i>Sma I</i> , <i>Bgl II</i>	
	pAcGHLT-C (8.7 kbp)	<i>polyhedrin</i>	ATG, GST, <i>BamH I</i> , 6x His tag-Protein kinase A site, Thrombin cleavage, <i>Nde I</i> , <i>Xho I</i> , <i>EcoR I</i> , <i>Stu I</i> , <i>Nco I</i> , <i>Sac I</i> , <i>Not I</i> , <i>Sse8387 I</i> , <i>Pst I</i> , <i>Kpn I</i> , <i>Sma I</i> , <i>Bgl II</i>	
	pAcGP67-A (9.8 kbp)	<i>polyhedrin</i>	ATG, GP64 signal sequence, <i>BamH I</i> , <i>Sma I</i> , <i>Xba I</i> , <i>EcoR I</i> , <i>Not I</i> , <i>Eag I</i> , <i>Pst I</i> , <i>Bgl II</i> , <i>Ppum I</i>	A, B and C represent three different reading frames and also contain slightly different restriction sites.
	pAcGP67-B (9.8 kbp)	<i>polyhedrin</i>	ATG, GP64 signal sequence, <i>BamH I</i> , <i>Sma I</i> , <i>Nco I</i> , <i>EcoR I</i> , <i>Not I</i> , <i>Eag I</i> , <i>Pst I</i> , <i>Bgl II</i>	
	pAcGP67-C (9.8 kbp)	<i>polyhedrin</i>	ATG, GP64 signal sequence, <i>BamH I</i> , <i>Sma I</i> , <i>Nco I</i> , <i>EcoR I</i> , <i>Not I</i> , <i>Eag I</i> , <i>Pst I</i> , <i>Bgl II</i> , <i>Ppum I</i>	
	pAcHLT-A (8.1 kbp)	<i>polyhedrin</i>	ATG, GST, <i>BamH I</i> , 6x His tag, Protein kinase A site-Thrombin cleavage, <i>Nde I</i> , <i>EcoR I</i> , <i>Stu I</i> , <i>Nco I</i> , <i>Sac I</i> , <i>Not I</i> , <i>Sse8387 I</i> , <i>Pst I</i> , <i>Kpn I</i> , <i>Sma I</i> , <i>Bgl II</i>	A, B and C represent three different reading frames and also contain slightly different restriction sites.
	pAcHLT-B (8.1 kbp)	<i>polyhedrin</i>	ATG, GST, <i>BamH I</i> , 6x His tag, Protein kinase A site, Thrombin cleavage- <i>Xho I</i> , <i>EcoR I</i> , <i>Stu I</i> , <i>Nco I</i> , <i>Sac I</i> , <i>Not I</i> , <i>Sse8387 I</i> , <i>Pst I</i> , <i>Kpn I</i> , <i>Sma I</i> , <i>Bgl II</i>	
	pAcHLT-C (8.1 kbp)	<i>polyhedrin</i>	ATG, GST, 6x His tag, Protein kinase A site, Thrombin cleavage site, <i>Nde I</i> , <i>Xho I</i> , <i>EcoR I</i> , <i>Stu I</i> , <i>Nco I</i> , <i>Sac I</i> , <i>Not I</i> , <i>Sse8387 I</i> , <i>Pst I</i> , <i>Kpn I</i> , <i>Sma I</i> , <i>Bgl II</i>	
	pBAC-1 (5.3 kbp)	<i>polyhedrin</i>	<i>BamH I</i> , <i>Stu I</i> , <i>EcoR I</i> , <i>Sac I</i> , <i>Hind III</i> , <i>Eag I</i> , <i>Not I</i> , <i>Ava I</i> , <i>Xho I</i> , 6x His tag, <i>Sty I</i> , <i>Avr II</i> , <i>Bpu 1102</i> , <i>Sph I</i>	

	pBacgus-1 (7.4 kbp)	<i>polyhedrin</i>	<i>Bam</i> HI, <i>Stu</i> I, <i>EcoR</i> I, <i>Sac</i> I, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Ava</i> I, <i>Xho</i> I, 6x His tag, <i>Sty</i> I, <i>Avr</i> II, <i>Bpu</i> 1102, <i>Sph</i> I	B-glucuronidase under p6.9 promoter control to monitor recombinant virus production.
	pBAC-2cp (5.4 kbp)	<i>polyhedrin</i>	<i>Nco</i> I, 6x His tag, <i>Sac</i> II, thrombin cleavage site, S tag, <i>PfIM</i> I, <i>Nhe</i> I, enterokinase cleavage site, LIC site, <i>Sma</i> I, <i>Srf</i> I, <i>BseR</i> I, <i>Stu</i> I, <i>BamH</i> I, <i>EcoR</i> I, <i>Sac</i> I, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Xho</i> I, 6x His tag, <i>Avr</i> II, <i>Bpu1102</i> I, <i>Sph</i> I	A Ligation-Independent Cloning (LIC) version of the vector is available for directional cloning PCR products.
	pBacgus-2cp (7.6 kbp)	<i>polyhedrin</i>	<i>Nco</i> I, 6x His tag, <i>Sac</i> II, thrombin cleavage site, S tag, <i>PfM</i> I, <i>Nhe</i> I, enterokinase cleavage site, LIC site, <i>Sma</i> I, <i>Srf</i> I, <i>BseR</i> I, <i>Stu</i> I, <i>BamH</i> I, <i>EcoR</i> I, <i>Sac</i> I, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Xho</i> I, 6x His tag, <i>Avr</i> II, <i>Bpu1102</i> I, <i>Sph</i> I	Ligation-Independent Cloning (LIC) version of the vector is available for directional cloning PCR products. B-glucuronidase under p6.9 promoter control to monitor recombinant virus production A.
	pBAC-3 (5.5 kbp)	<i>polyhedrin</i>	Gp64 signal peptide, <i>Nco</i> I, 6 x His tag, <i>Sac</i> II, thrombin cleavage site, S tag, <i>PfIM</i> I, <i>Nhe</i> I, enterokinase cleavage site, <i>Sma</i> I, <i>Srf</i> I, <i>BseR</i> I, <i>Stu</i> I, <i>BamH</i> I, <i>EcoR</i> I, <i>Sac</i> I, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Xho</i> I, 6 x His tag, <i>Avr</i> II, <i>Bpu1102</i> I, <i>Sph</i> I	
	pBAC-5 (5.5 kbp)	<i>gp64</i>	<i>Nco</i> I, 6x His tag, <i>Sac</i> II, Thrombin cleavage site, S tag, <i>PfIM</i> I*, <i>Nhe</i> I, enterokinase cleavage site, <i>Sma</i> I, <i>Srf</i> I, <i>BseR</i> I, <i>Stu</i> I, <i>BamH</i> I, <i>EcoR</i> I, <i>Sac</i> I, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Xho</i> I, 6x His tag, <i>Avr</i> II, <i>Bpu1102</i> I, <i>Sph</i> I	Has early/late gp64 promoter but no signal peptide coding region.

\* within S tag

	pBACgus-5 (7.7 kbp)	<i>gp64</i>	Nco I, 6x His tag, Sac II, Thrombin cleavage site, S tag, PflM I*, Nhe I, enterokinase cleavage site, Sma I, Srf I, BseR I, Stu I, BamH I, EcoR I, Sac I, Hind III, Eag I, Not I, Xho I, 6x His tag, Avr II, Bpu1102 I, Sph I  * within S tag	B-glucuronidase under p6.9 promoter control to monitor recombinant virus production.  Has early/late gp64 promoter but no signal peptide coding region.
	pBAC-6 (5.6 kbp)	<i>gp64</i>	GP64 signal sequence, Nco I, 6x His tag, Sac II, Thrombin cleavage site, S tag, PflM I*, Nhe I, enterokinase cleavage site, Sma I, Srf I, BseR I, Stu I, BamH I, EcoR I, Sac I, Hind III, Eag I, Not I, Xho I, 6x His tag, Avr II, Bpu1102 I, Sph I.  * within S tag	Has early/late gp64 promoter and signal peptide coding region.
	pAcSecG2T (8.6 kbp)	<i>polyhedrin</i>	ATG, GP64 signal sequence, GST, Thrombin cleavage site, BamH I, Sma I, EcoR I	
	pAcSG2 (5.5 kbp)	<i>polyhedrin</i>	Xho I, EcoR I, Stu I, Nco I, ATG, Sac I, Not I, Eag I, Sse8387 I, Pst I, Kpn I, Sma I, Bgl II	ATG codon for fusion proteins.
<b>Multiple Promoter (<i>polyhedrin</i>)</b>	pAcUW21 (9.2 kbp)	<i>polyhedrin</i>  <i>p10</i>	See note  <i>Pac</i> I, <i>Bgl</i> II, <i>EcoR</i> I	Polyhedrin gene intact; foreign gene expression from p10 promoter.
	pAcUW51 (5.8 kbp)	<i>polyhedrin</i>  <i>p10</i>	<i>Bam</i> H I  <i>Bgl</i> II, <i>EcoR</i> I	Dual expression vector from polyhedrin and p10 promoters.

	pAcAB3 (10.1 kbp)	<i>p10</i> <i>polyhedrin</i> <i>p10</i>	<i>Sma</i> I, <i>Bam</i> H I <i>Xba</i> I, <i>Stu</i> I <i>Bgl</i> II, <i>Esp</i> I	Triple expression vector from polyhedrin and 2 x p10 promoters.
	pAcDB3 (6.0 kbp)	<i>p10</i> <i>polyhedrin</i> <i>p10</i>	<i>Sma</i> I, <i>Bam</i> H I <i>Xba</i> I, <i>Stu</i> I <i>Bgl</i> II, <i>EcoR</i> I	Triple expression vector from polyhedrin and 2 x p10 promoters. Smaller version of pAcAB3.
	pAcAB4 (10.2 kbp)	<i>polyhedrin</i> <i>p10</i> <i>polyhedron</i> <i>p10</i>	<i>Bam</i> H I <i>Sma</i> I <i>Xba</i> I, <i>Stu</i> I <i>Bgl</i> II, <i>EcoR</i> I, <i>Esp</i> I	Quadruple expression vector from 2 x polyhedrin and 2x p10 promoters.
	pBAC4x-1 (5.9 kbp)	<i>p10</i> <i>polyhedrin</i> <i>p10</i> <i>polyhedrin</i>	<i>Bgl</i> II, <i>EcoR</i> I, <i>Bsu</i> 36 I <i>Xba</i> I, <i>Stu</i> I <i>Sma</i> I, <i>Spe</i> I <i>Bam</i> H I, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Xho</i> I 6x His tag, <i>Sty</i> I, <i>Bpu</i> 1102 I, <i>Sph</i> I	Quadruple expression vector from 2 x polyhedrin and 2 x p10 promoters.

	pBACgus4x-1 (8.1 kbp)	<i>p10</i> <i>polyhedrin</i> <i>p10</i> <i>polyhedrin</i>	<i>Bgl</i> II, <i>EcoR</i> I, <i>Bsu36</i> I  <i>Xba</i> I, <i>Stu</i> I  <i>Sma</i> I, <i>Spe</i> I  <i>BamH</i> I, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Xho</i> I 6x His tag, <i>Sty</i> I, <i>Bpu1102</i> I, <i>Sph</i> I	Quadruple expression vector from 2 x polyhedrin and 2x p10 promoters. B-glucuronidase under p6.9 promoter control to monitor recombinant virus production.
	pBACgus-6 (7.7 kbp)	<i>gp64</i>	GP64 signal sequence, <i>Nco</i> I, 6x His tag, <i>Sac</i> II, Thrombin cleavage site, <i>S</i> tag, <i>PfIM</i> I*, <i>Nhe</i> I, enterokinase cleavage site, <i>Sma</i> I, <i>Srf</i> I, <i>BseR</i> I, <i>Stu</i> I, <i>BamH</i> I, <i>EcoR</i> I, <i>Sac</i> I, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Xho</i> I, 6x His tag, <i>Avr</i> II, <i>Bpu1102</i> I, <i>Sph</i> I  *within <i>S</i> tag	Has early/late gp64 promoter and signal peptide coding region. B-glucuronidase under p6.9 promoter control to monitor recombinant virus production.
	pBACsurf-1 (9.4 kbp)	<i>polyhedrin</i>	<i>Spe</i> I, gp64 signal sequence, <i>Pst</i> I, <i>Kpn</i> I, <i>Sma</i> I, gp64 coding region	Designed for incorporating target proteins on the virion surface by utilizing gp64 signal sequence and membrane anchor region.
	pTriEx-3 (5.1 kbp)	<i>CMV</i> immediate/early <i>T7lac</i> <i>p10</i>	<i>Nco</i> I, <i>EcoR</i> V, <i>Sma</i> I, <i>Ecl136</i> II, <i>Sac</i> I, <i>BamH</i> I, <i>EcoR</i> I, <i>Bgl</i> II, <i>BssH</i> II, <i>Asc</i> I, <i>Pst</i> I, <i>Sse8387</i> I, <i>Kpn</i> I, <i>PinA</i> I, <i>Nsp</i> V, <i>Hind</i> III, <i>Eag</i> I, <i>Not</i> I, <i>Pvu</i> II, <i>Bst1107</i> I, <i>Pml</i> I, <i>Xho</i> I, <i>Dra</i> III, <i>Bsu36</i> I	Suitable for the expression of a single recombinant plasmid in <i>E.coli</i> , insect and vertebrate systems by utilising one of three promoters ( <i>T7lac</i> , <i>p10</i> and <i>CMV</i> respectively).
Single Promoter ( <i>Basic/p6.9</i> )	pAcMP2 (9.8 kbp)	<i>Basic</i> ( <i>p6.9</i> )	<i>Pst</i> I, <i>Not</i> I, <i>Eag</i> I, <i>EcoR</i> I, <i>Xba</i> I, <i>BamH</i> I	<i>p6.9</i> promoter provides late gene expression.
	pAcMP3 (9.8 kbp)		<i>BamH</i> I, <i>Xba</i> I, <i>EcoR</i> I, <i>Not</i> I, <i>Eag</i> I, <i>Pst</i> I, <i>Bgl</i> II	
Single Promoter ( <i>p10</i> )	pAcUW1 (4.6 kbp)	<i>p10</i>	<i>Bgl</i> II, <i>Hind</i> III	

<b>Multiple Promoter (<i>p10</i>)</b>	<b>pAcUW42 (7.1 kbp)</b>	<i>p10</i> <i>polyhedrin</i>	<i>Bgl</i> II, <i>Pst</i> I, <i>Not</i> I, <i>Xba</i> I, <i>Kpn</i> I, <i>Sma</i> I  <i>Bam</i> H I	
	<b>pAcUW43 (7.1 kbp)</b>	<i>p10</i> <i>polyhedrin</i>	<i>Sma</i> I, <i>Kpn</i> I, <i>Xba</i> I, <i>Not</i> I, <i>Pst</i> I, <i>Bgl</i> II  <i>Bam</i> H I	